



A contribution to the synsedimentary versus epigenetic origin of the Cu mineralizations hosted by terminal Neoproterozoic to Cambrian formations of the Bou Azzer–El Graara inlier: New insights from the Jbel Laassel deposit (Anti Atlas, Morocco)

Hugo Bourque, Luc Barbanson, Stanislas Sizaret, Yannick Branquet, Claire Ramboz, A. Ennaciri, M. El Ghorfi, L. Badra

► To cite this version:

Hugo Bourque, Luc Barbanson, Stanislas Sizaret, Yannick Branquet, Claire Ramboz, et al.. A contribution to the synsedimentary versus epigenetic origin of the Cu mineralizations hosted by terminal Neoproterozoic to Cambrian formations of the Bou Azzer–El Graara inlier: New insights from the Jbel Laassel deposit (Anti Atlas, Morocco). *Journal of African Earth Sciences*, 2015, 107, pp.108-118. 10.1016/j.jafrearsci.2015.04.005 . insu-01145561

HAL Id: insu-01145561

<https://hal-insu.archives-ouvertes.fr/insu-01145561>

Submitted on 12 Jun 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

A contribution to the synsedimentary versus epigenic origin of the Cu mineralizations hosted by terminal Neoproterozoic to Cambrian formations of the Bou Azzer – El Graara inlier: new insights from the Jbel Laassel deposit (Anti Atlas, Morocco)

H. Bourque, L. Barbanson, S. Sizaret, Y. Branquet, C. Ramboz, A. Ennaciri, M. El Ghorfi, L. Badra

PII: S1464-343X(15)00078-3

DOI: <http://dx.doi.org/10.1016/j.jafrearsci.2015.04.005>

Reference: AES 2252

To appear in: *African Earth Sciences*

Received Date: 6 October 2014

Revised Date: 3 April 2015

Accepted Date: 4 April 2015



Please cite this article as: Bourque, H., Barbanson, L., Sizaret, S., Branquet, Y., Ramboz, C., Ennaciri, A., El Ghorfi, M., Badra, L., A contribution to the synsedimentary versus epigenic origin of the Cu mineralizations hosted by terminal Neoproterozoic to Cambrian formations of the Bou Azzer – El Graara inlier: new insights from the Jbel Laassel deposit (Anti Atlas, Morocco), *African Earth Sciences* (2015), doi: <http://dx.doi.org/10.1016/j.jafrearsci.2015.04.005>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**A contribution to the synsedimentary versus epigenic origin of the Cu
mineralizations hosted by terminal Neoproterozoic to Cambrian formations
of the Bou Azzer – El Graara inlier: new insights from the Jbel Laassel
deposit (Anti Atlas, Morocco).**

H. Bourque ^{(1)*}, L. Barbanson ⁽¹⁾, S. Sizaret ⁽¹⁾, Y. Branquet ⁽¹⁾, C. Ramboz ⁽¹⁾, A. Ennaciri ⁽²⁾, M. El
Ghorfi ⁽³⁾ and L. Badra ⁽⁴⁾

⁽¹⁾ *Institut des Sciences de la Terre d'Orléans (ISTO), UMR 7327-CNRS/Université d'Orléans/BRGM, Orléans,
France*

⁽²⁾ *Managem Group, Casablanca, Morocco*

⁽³⁾ *Faculté des Sciences et Techniques – Guéliz, Bd. Abdelkrim El Khattabi - B.P. 549– Marrakech, Morocco*

⁽⁴⁾ *Université Moulay Ismaïl, Meknès, Morocco*

* Correspondence to:

hugo.bourque@cnrs-orleans.fr

Institut des Sciences de la Terre d'Orléans

UMR 7327-CNRS/Université d'Orléans

1A rue de la Ferrollerie

45071 Orléans Cedex 2

France

T : +33 (0)2 38 49 27 64

21

22 **Abstract**

23 The Neoproterozoic to Cambrian formations that compose the cover of the Bou Azzer-El
24 Graara inlier, host a great number of Copper occurrence whose origin is largely discussed. To bring
25 some light to this debate, structural, petrographic and geochemical observations were performed on
26 the copper deposit of Jbel Laassel. This deposit, located at the extreme ESE of the Bou Azzer –El
27 Graara inlier, is mined since 2012. At the district scale, the ore bodies localize in a folding band that
28 extends along a NE-SW direction. At macroscopic, microscopic and scanning electron microscope
29 scales the mineralization appears as banding veins, with locally cockade breccia and comb quartz
30 textures. From the macroscopic scale to the scale of the scanning electron microscope, all these
31 mineralized textures are connected there between forming a stockwork with an auto-similar structure
32 in the range of used scales of observation. At the district scale, this stockwork is preferentially located
33 in the anticlinal hinges of the folding band. Principal component analyses of geochemical database
34 enable to distinguish several groups of chemical elements, each of these groups corresponding to the
35 different lithologies and to the copper mineralization. This last group doesn't show any correlation
36 with the distinguished lithological groups. All these observations bring new arguments to an
37 epigenetic origin for the copper mineralization of the Jbel Laassel deposit, with a formation
38 contemporary or posterior with the folding band development attributed to Variscan deformation.

39

40 **Keywords:** Anti-Atlas, Cu-mineralization, folding band, stockwork, epigenetic.

41 **1. Introduction**

42 More than 200 Copper mineralizations are known over a large part of the Neoproterozoic to
43 Cambrian cover in the Anti-Atlas (fig. 1A) (Bouchta et al., 1977). They are localized at different
44 stratigraphic levels within the cover and present different characteristics. Their origin remains in most

45 cases currently poorly understood and there is no general model for this copper mineralization. Several
46 genetic interpretations have been proposed on specific deposits: (i) based on textural and petrological
47 observations, Leblanc (1986) suggests that Alous mineralization crystallized during the cooling of an
48 ignimbrite; (ii) For the Cu-occurrences of Tizert, Talat N'Ouamane, Tizirt and Amadou, Pouit
49 (1966), Bouchta & al (1977) and Skacel (1993) consider these mineralizations generated through a
50 synsedimentary process arguing of a strong paleogeographic control of the ores. Moreover, the Cu-
51 mineralizations hosted in the Neoproterozoic to Cambrian cover in the Anti-Atlas can be differentiate
52 by their morphologies as veins, dissemination or stratiform bodies, without relationship have been
53 established between these morphologies yet (Pouit, 1966; Skacel, 1993). As the result, the syngenetic
54 or epigenetic nature of that mineralization remains still undetermined (Pouit, 1966). New data and
55 evidences are then necessary. The Jbel Laassel deposit is one of these numerous cover-hosted copper
56 occurrences known in the Anti-Atlas terminal Neoproterozoic to Cambrian cover, often called
57 Adoudounian cover (Pouit, 1966; Bouchta et al., 1977; Benssaou & Hamouni, 1999). Since the 60's,
58 various genetic models were proposed by several geologists hired by Managem Goup for the copper
59 mineralization at Jbel Laassel site. Maacha et al. (2011) resumed these different models: in 1964 the
60 mineralization was attributed to a porphyry copper type, in 1967 it was interpreted as a stratiform
61 synsedimentary deposit, with a re-concentration stage associated to a Jurassic doleritic intrusion, in
62 1978 an epigenetic origin was proposed for this mineralization with a mainly vein-shaped texture and
63 a close association with barite. Some authors evoked a synsedimentary genesis for the Jbel Laassel
64 deposit, where mineralization is controlled by the basement paleogeography and were locally
65 remobilized by a tectonic event (Pouit, 1966; Bouchta et al., 1977). According to Skacel (1993),
66 copper should be synchronous with sedimentation; its precipitation being regulated by the redox
67 conditions, themselves under the control on the deposition environment.

68 In 1984, the MANAGEM Group discovered the potential of the Jbel Laassel Cu-
69 mineralizations. Different preliminary estimations were performed until 2006 (Maacha et al., 2011). In
70 2010 Managem decided to lead a core drill campaign to estimate the feasibility of this deposit. This
71 work resulted in an estimation of 7.5 million tons at 1% Cu (Maacha et al., 2011) and the exploitation

began in 2012 with SOMIFER as operator. In this article, based on MANAGEM Group pre-exploitation targeting, we report the main results of an original study performed on Jbel Laassel Cu-mineralization. This work consists of: (i) a structural analysis of ore bodies, (ii) a mineralogical study of samples collected both on outcrops and drill cores and (iii) a statistical analysis of the first chemical analyses carried out in this deposit. These new data bring out new insights on the debated, epigenic versus syngenetic origin of the Jbel Laassel deposit.

2. Geological setting

2.1. The Bou Azzer El Graara inlier

The Bou Azzer El Graara inlier is one of a series of Proterozoic windows oriented NW-SE that expose Panafrican formations in the central part of the Anti-Atlas (Choubert, 1947). The Proterozoic basement is mainly composed by a dismembered ophiolitic sequence and arc fragments (Leblanc, 1975; Saquaque et al., 1989) (fig. 1B). These formations are unconformably overlain by a thick Neoproterozoic to Cambrian volcano-sedimentary cover (Soulaimani et al., 2014). This cover can be divided into three formations from bottom to top: (1) The Tiddiline Formation (~750 to 650 Ma) attributed to the “Saghro Group” (Thomas et al. 2004), composed of clastic, volcanoclastic and volcanic series, rests mainly uncomfortably on the Panafrican substratum; (2) The Ouarzazate Group (~610 to 550 Ma), composed of a volcano sedimentary complex, rests in angular unconformity on the Tidiline Formation; (3) Terminal Neoproterozoic to Cambrian Formations consisting of detrital and carbonated series (Soulaimani et al., 2013). These Terminal Neoproterozoic to Cambrian Formations, varying in age from terminal Neoproterozoic to the middle Cambrian, are associated with a major marine transgression toward the Southeast and can be subdivide into two groups: the Taroudannt Group and the Tata Group (fig. 2). During the Late Paleozoic compressional event, the Panafrican structures of the basement were reactivated along the inlier's borders. This results in box-shaped folds distributed throughout the Bou Azzer-El Graara area and by large open synclines of Cambrian rocks (Soulaimani & Burkhard, 2008). Upright detachment folds, from meters to decameters in scale, are

97 frequent in the lower Cambrian rocks and they exhibit a dominant NW-SE trend with subordinate NE-
 98 SW structures (Soulaimani & Burkhard, 2008).

99

100 **2.2. Lithostratigraphy of the Jbel Laassel Cu-deposit**

101 The Jbel Laassel deposit is hosted in the Lower Cambrian part of the Neoproterozoic to
 102 Cambrian volcano-sedimentary cover, at the extreme ESE part of the Bou Azzer-El Graara inlier, 30
 103 kilometers NE of the Bleida mine (fig. 1B). The whole of this cover is often call Adoudounian cover,
 104 it is a local appellation, and on the other hand the term Adoudou correspond to the name of a precise
 105 stratigraphic formation (Soulaimani et al, 2013). The Lower Cambrian is represented in the studied
 106 site by the Adoudou and Tikirt Formations, both belonging to the Taroudann Group, whereas the
 107 Igoudine, Amouslek, Issafene and Tazlaft Formations correspond to the Tata Group (fig. 2)
 108 (Soulaimani et al., 2013). All these formations are interlayered locally with volcanic flows dated at
 109 534 ± 10 Ma (U/Pb on zircon by Ducrot & Lancelot, 1977) related to the Jbel Boho-type volcanism
 110 (Alvaro et al., 2006) (fig. 1 and 2).

111 **2.2.1 The Taroudannt Group**

112 The Adoudou Formation (Choubert, 1952) is represented in the area study by a basal
 113 sedimentary breccia with a thickness of 5 to 100 meter and by an alternation of dolostones and red or
 114 white clay's siltstones with a total thickness varying from 150 to 250 meters. Dolostones frequently
 115 present an intensive secondary silicification (Soulaimani et al., 2013). The Tikirt Formation (250 to
 116 300 meters thick) rests in conformity on the Adoudou Formation; it is composed essentially by
 117 sandstones frequently interlayered by centimetric clayed siltstones. The age of this formation is
 118 unknown; but the Taliwine Formation, a westward stratigraphic equivalent level has been dated by
 119 U/Pb on zircon from Early Cambrian volcanic horizons at 521 ± 7 Ma and 522 ± 2 Ma (Compston et
 120 al., 1992; Landing, 1998; Maloof et al., 2005). On the Jbel Laassel site and at regional scale, Tikirt
 121 sandstones end with a continuous level of red claystones 8 to 15 meters thick (fig. 2).

2.2.2 The Tata Group

This Group includes, from oldest to youngest, the formations of Igoudine, Amouslek, Issafène and Tazlaft. The age of the Tata Group varies from the Tomotien (in the lower Cambrian) at its base, to the Middle Cambrian at its top. (Soulaïmani et al., 2013). The Igoudine Formation (50 to 60 meters thick) is made up of dolostones with microbialite interlayered with clayed siltstones or siltstones. The Amouslek Formation appears as an alternation of stromatolitic dolostones with heterolithic facies of clayed siltstones, for a total thickness of 150 to 180 meters. The Issafène Formation is composed by red claystones and thin beds of sandstones, and it is 30 to 40 m thick (Soulaïmani et al., 2013). The Tazlaft Formation, 90 to 100 meters thick, is composed of sandstones with oblique cross bedding and mega-ripples marks (Soulaïmani et al., 2013).

3. Methodology and Analytical procedure

In order to decipher the complex pattern of geometries and structures, a structural study has been performed in the field coupled with high density drill control. As a result, well constrained and high resolution maps and cross-sections have been constructed (fig. 3 to 5).

During the field work, 10 samples have been collected from outcrops and 55 samples have been collected on 9 core drills. 60 polished thin sections were prepared from these samples. They have been first observed using a Leica DMRX petrographic microscope (transmitted and reflected light modes). Complementary observations and analyses were carried out using JSM-6400 JEOL Scanning Electron Microscope (SEM) at ISTO. Polished thin-sections have been first coated by a thin carbon layer. Acceleration voltage and beam current were 20 kV and 8 nA, respectively. IdFix Software package was used for data processing. Back-scattered electrons (BSE) imaging mode was used to reveal the composition variations at microscopic scale, whereas the texture of clayey material was examined with secondary electrons (SE) imaging mode. The SEM system is coupled to an Energy-Dispersive X-ray spectrometer (EDS) to make qualitative determinations of the mineral composition.

The Managem Group performed analyses on 23 Reverse Circulation drilling's samples collected at different depths for the following elements: Cu, CuOx, SiO₂, Al₂O₃, Fe₂O₃, CaO, MgO, K₂O, MnO, TiO₂, P₂O₅, As, B, Ba, Be, Bi, Cd, Co, Cr, Ge, Li, Mo, Nb, Ni, Pb, Sb, Se, Sn, Sr, W, Y, Zn, Ag and loss on ignition (LOI). "Cu" corresponds to the total copper content and CuOx corresponds to the non-sulphide copper content. For the major elements analysis, 0.5 g of the sample is crushed at less than 100µm and it is dissolved by fusion at 500°C during 45 minutes with 2.5 g of sodium peroxide in a zirconium crucible. The melt mixing is dissolve with 100 ml of hydrochloric acid (28% HCl) and the solution is analyzed using an ICP-AES ULTIMA 2C using the Jobin Yvon-HORIBA device. For the other elements, 0.25 g of the sample is dissolved by acid attack (50%HCl and 50% HNO₃) microwave-assisted during 45 minutes at 220°C. The solution is then analyzed by ICP-MS Thermo X'Serie 2.

4. Results

4.1. Structural study

The whole sequence undergoes at least one deformation stage. Two types of folds are present. The first family corresponds to upright folds characterized by sub-horizontal to slightly NW dipping axes with a NW-SE to NNW-SSE orientation (fig. 3b). They are marked by an axial planar cleavage particularly developed in the fold hinge zone. Those NW-trending folds are concentrated in a band, named herebelow "folding band", which displays a width of 150 to 200 meters and is oriented N145° (fig. 3 and 4). This "folding band" is developed along a N150E-trending vertical fault, folds being localized in both sides of the fault. The second fold family shows moderately inclined fold's axes to the NE, oriented NE-SW (fig. 3). The orientation of NE-trending folds is parallel to the direction (around N30°) of a SE-vergent thrust that outcrops through the entire area (fig. 3). Therefore, at large scale, the Jbel Laassel area and its ore body appears to be located within a large NE-trending synclinal developed in the thrust footwall (figure 3). The relations between the two types of folds could not be observed in the field. In the thrust hanging wall, a similar vertical fault is found northeastward (fig. 3) with sub-parallel folds and mineralizations along the northwestern prolongation (out of fig. 3).

Consequently, we infer that both hanging- and footwall vertical fault segments are part of the same vertical fault offset by the late thrust. As the fault is vertical, the thrust might have right lateral oblique slip component of about 300 meters.

4.2. Jbel Laassel ore bodies

4.2.1. Geometries and orientation

The vertical and lateral extent of ore bodies was delimited by surface cartography and drills. Part of the outcrops are covered by superficial alteration formations and mining waste. Ore bodies extend 150 to 250 meters in width (fig. 4), with a maximal longitudinal extension of 400 meters (fig. 3) for an average thickness of around 100 meters (fig. 4 and 6). Mineralization is hosted in Igoudine and Amouslek Formations and more especially in dolostones and siltstones beds, rarely in lavas or claystones beds. It is noteworthy that ore bodies are distributed within and along the “folding band” with an orientation NE-SW i.e., they show the same preferential orientation as that of the folds (fig. 3). Moreover, on a NE-SW cross section (fig. 4), the ore bodies show thickness variation: in the hinge of anticlinal the thickness is maximal whereas in the hinge of synclinal the thickness is minimal (fig. 4). Similarly, the vertical fault strongly controls the ore bodies distribution with a nearly barren southwestern compartment contrasting with a rich folded northeastern one (fig. 4). On a NW-SE cross section, the thickness of the ore bodies is constant and does not show significant variation (fig. 5).

4.2.2 Mineralogical composition and paragenetic succession

Without distinction between primary and secondary origin, the copper mineralization is composed of chalcocite, bornite, chalcopyrite, covellite, digenite, malachite, chrysocolla, tenorite, native copper and cuprite, associated with quartz, dolomite and calcite as gangue mineral. These gangue minerals in veinlets or voids systematically show, a banding texture. The observed growth direction indicates a centripetal quartz, dolomite and finally calcite (fig. 6A). This gangue chronology

is a good landmark to determine the copper mineral succession. Bornite appears as minute grains (less than 20 μm) in the quartz and dolomite ribbons, equally distributed between these two minerals. Frequently, lamella exsolutions of chalcopyrite are observed in bornite grains. Chalcocite occurs in ribbon between the bands of dolomite and calcite. Chalcocite grains mostly appear with an anhedral shape without internal structure. Sometimes this mineral displays a hexagonal cleavage (fig. 6B and 6C), suggesting, in this case, a primary origin at above 103°C (Ramdhor, 1969). This generation of chalcocite is named CC1. Beside, chalcocite exists in association with covellite, digenite and chalcopyrite, in replacement around the bornite, in this case chalcocite, covellite and digenite can be interpreted as cementation assemblage. This chalcocite is named CC2. Malachite replaces chalcocite and dolomite and may be replaced by chrysocolla. Malachite also replaces cuprite but sometimes the opposite situation is observed. Veinlet fills only by malachite are been observed, they cut all the other rock components and structures except chrysocolla.

All these relations between minerals allow to discriminate a primary origin represented by: quartz, bornite, chalcopyrite, dolomite, chalcocite (CC1) and calcite, and a secondary origin represented by: covellite, digenite, chalcocite (CC2), malachite, chrysocolla, tenorite, native copper and cuprite (fig. 7). Based on intersections relations between the different mineral phases, several episodes of fissuring have been observed (fig. 7).

4.2.3 Multiscale observations of the ore texture

At macroscopic scale, on core samples, two textural types of mineralization have been distinguished. The most frequent is a stockwork with veins/veinlets parallel or oblique to the stratification plane (fig. 8). Parallel-bedding veins are still connected to cross-bedding veins (fig. 8C). The second type corresponds to disseminations within the rock ground mass. Locally it is spatially related to the stockwork (fig. 8D). When it is well develop, veins of the stockwork can show breccia texture with angular fragments and in situ fragmentation texture without significant rotation of the fragment that is characteristic of fluid-assisted brecciation (Jébrak, 1997) (fig. 8A). At the microscopic scale, both types of mineralization appear composed by the same ore and gangue minerals. Textural

and chronological relationships being also similar, both mineralization types have the same paragenetic sequence (fig. 6, 7 and 8). Veins can also locally show a cockade breccia and quartz with comb texture (fig. 6F and 6G). At this scale, disseminated grains of copper-bearing minerals, down to 50 micrometers in size, are still observable (fig. 6E). Noteworthy, they frequently show a close spatial relationship with the veinlet of the stockwork (fig. 6E). At the scanning electron microscope scale, the disseminated mineralization appears as micro-voids or geodes that display the same mineralogical content and the same paragenetic succession as that observed at higher observation scales (i.e. with quartz or dolomite at the wallrock and chalcocite at the center) (fig. 9).

4.2.4 Mineralogical distribution at the deposit scale

Using the data collected on 9 core drills and on surface mapping, the Cu-bearing mineral abundance is plotted in cross-sections (fig. 4 and 5). The mineralogical abundances are displayed in the form of spider diagram, with, at the top, the secondary Cu-bearing "oxydized" minerals (malachite, chrysocolla and cuprite) and at the bottom the primary or cementation Cu-bearing minerals (chalcocite, bornite and chalcopyrite). The mineralogical abundances are evaluated through microscopic observations of polished thin sections of mineralized cores. The content of each mineral is evaluated visually using an abundance chart (Dutro et al., 1989). Results are expressed in the corresponding axis of the spider diagram in a scale of 0 to 100% with steps of 20%.

Chalcocite and malachite are the most abundant minerals in the Jbel Laassel deposit (fig. 4 and 5). According to our observations, no obvious zonation appears between primary sulfides and "oxidized" mineralization.

4.3. Principal Component Analysis (PCA) of chemical data

Two multi-element databases were available for PCA: the larger one, does not include Cu analyses, while the second one includes Cu analyses for a smaller number of samples. PCA analysis was performed on both sets separately using varimax criteria.

4.3.1. Dataset without copper measurements

The first results on database without copper measurements (228 analyses) show that two factors F1 and F2 explain 42.06 % of the total variability (fig. 10A). Three groups of elements discriminated in the F1-F2 plane represented by the CaO, MgO and MnO cluster (group 1) discriminates the carbonate matrix, the second SiO₂, Al₂O₃, K₂O and B clusters represents the clay sedimentary component (group 2) whereas the last Co, P₂O₅, Fe₂O₃, TiO₂ and Pb cluster (group 3) probably marks the lava component (fig. 10B). Thus, referring to the F1 F2 chemical space (fig. 10C), the analyzed dolostones consists of alternation of siltstones and dolostone layers, with interstratified lavas in both of them. It appears that the three groups of variables can be interpreted in terms of lithology: the groups 1, 2 and 3, corresponding respectively to dolostones, siltstones and lava formations (fig. 10C).

4.3.2 Dataset including copper measurements

This data set comprises 72 analyses. For the Cu-mineralized data set (Cu % and CuOx %), the four first factors account for 80% of the total variance. Beyond factor 4 the explained variance decreases sharply. The plane F1-F2 represents 48.29% of the total variance (28.27% for F1 and 20.01 for F2). The three petrographic groups previously identified in the F1-F2 plane are well discriminated by the same element clusters (fig. 10D). This representation of the variables suggests that copper (total or oxidized) is not linked to any of the three types of lithology. Indeed, copper is independent of sedimentary dolostones and siltstones, and it is anticorrelated with the lavas (fig. 10D). Moreover, an examination of the F1-F2 plane, suggests that copper could be associated with barium, but the correlation coefficient between total copper and barium is -0.05 and that of barium with oxidized copper is 0.03. This is due to the fact that the total copper, oxidized copper and barium are badly represented in the plane F1 - F2, as shown in the diagram in figure 10D and by the values of the weighting coefficients between the original variables and factors. For total copper, oxidized copper and barium, these values are -0.19, -0.23 and -0.16 with respect to F1 and -0.30, -0.38 and -0.38 relative to F2. The factors where copper (total and oxidized) is best represented are F3 and F6. The

weighting coefficients between the copper, the oxidized copper and the factors are 0.54 and 0.42 with respect to F3 and 0.59 and 0.63 relative to F6, respectively. The variables representation in the plane F3-F6 (17.37% of the total variance, 12.55% for F3 and 4.82% for F6) shows that the variations of the total copper and of the oxidized copper are not associated with changes in the others variables values (fig. 10E). Therefore, the PCA highlights the independence of copper (total and oxidized) relative to the others variables present in the available database, especially that corresponding to lithology.

5. Discussion

In the Jbel Laassel copper deposit, at a macroscopic scale, the mineralization is present as a stockwork and disseminations. The stockwork is composed by veins filled with a primary paragenesis composed by: quartz, bornite, chalcopyrite, dolomite, chalcocite, calcite and a secondary paragenesis composed by: covellite, digenite, malachite, chrysocolla, tenorite, cuprite and native copper. The veins of this stockwork crosscut the stratification plane or are parallel to it. Both vein families always present the same mineralogy, texture and paragenetic evolution. These observations suggest that, with respect to the primary sulfide paragenesis, all the stockwork veins are coeval and formed during the same mineralizing event. SEM observations reveal that disseminations correspond to microcracks and cavities filled with the same mineralogical content and the same textural characteristics that of stockwork veins; moreover the disseminated Cu-bearing mineral grains are directly “connected” to the stockwork veins. Thus, all the primary mineralization, from the macroscopic scale to the scanning electron microscope scale, is related to the stockwork and deposited during a unique hydrothermal event of fluid assisted fracturation of the host sedimentary rocks. All these observations favor an epigenetic origin of the Jbel Laassel mineralization. In addition, the principal component analysis shows that copper does not have affinities with any of the lithological groups present in the area. This result is also consistent with an epigenetic origin of the mineralization. The limits of the ore body correspond to the limits of the “folding band” (fig. 3). In the cross-section perpendicular to the main extension of the folding band (fig. 4), the ore body is concentrated in the anticline hinges and his thickness decreases in syncline hinges; whereas in cross section parallel to the main extension of the folding axis, the ore body is continuous and does not show noticeable thickness variations (fig. 5).

Consequently the mineralized stockwork is clearly controlled by hectometric folds, which are, with the Igoudine and Amouslek Formations, the trap of the mineralization. Available structural and fault analysis data do not allow us to decipher clearly the control exerted by the vertical faults on NW-trending mineralized folds. However, the structural pattern associated with the "folding band" suggest folding developed within the cover above fault involving basement through drap folding and/or flower structure mechanisms (Sylvester, 1988; Fossen et al., 2013).

Soulaimani & Burkhard (2008) describes variscan kink bands with a rough cleavage in the vicinity of Panafrican structures in the basement of the Bou Azzer El Graara inlier. These authors interpret these folds in terms of the reactivation of Panafrican structures during the late Paleozoic compression. In the terminal Neoproterozoic to Cambrian's cover, folds are currently attributed to this Paleozoic compression, controlled by the movement of inherited basement structures (Leblanc, 1972; Soulaimani, 1997; Faïk et al, 2002; Soulaimani & Burkhard, 2008). It results in Bou Azzer-El Graara inlier, metric to decametric folds, have NW-SE trending axes exhibit with a subordinate NE-SW orientation (Soulaimani & Burkhard, 2008). The folds in the "folding band" of Jbel Laassel display the same characteristics as the ones given by Soulaimani & Burkhard (2008), for the Variscan folds, i.e. metric to decametric folds and NW-SE trending axes with a subordinate NE-SW orientation. In such conditions, the age of the mineralizing event could probably be contemporary to the deformation or younger. It could be possible that the intersection between the "folding band" and the NE-SW synclinal took a role in the trap of mineralization formation. This "folding band" is associated with a major NW-SE fault with a kilometers extension. This fault probably acted as a drain for fluids that formed the mineralization of the "folding band" in the Jbel Laassel area. Anticlinal hinges of the "folding band" could be a trap for hydrothermal discharges with a fluid-assisted fracturing focused in these hinges. The major NW-SE fault could be in direct relation with the basement, as reported by Soulaimani & Burkhard (2008) for the others Variscan faults in the terminal Neoproterozoic to Cambrian cover. This fault could thus emphasize a basement influence at the origin of the Cu mineralization of Jbel Laassel. We propose that Cu mineralization hosted in the terminal Neoproterozoic to Cambrian cover are linked to Variscan reactivation of inherited basement structures

and Variscan faults that drained fluids which precipitated its copper content in the anticlinal hinge of the Jbel Laassel "folding band".

9. Conclusion

The copper mineralization in Jbel Laassel area is epigenetic and is controlled and coeval with decametric folding, whose axes exhibit a NW-SE trend in the Igoudine and Amouslek Formations. These folds are concentrated in a 150 to 200 m wide band, probably formed during the Variscan compression. Consequently the mineralizing event took place during this period or later. The mineralization appears as a stockwork, displaying the same characteristics (texture and mineral composition) from the macroscopic scale to the scale of the scanning electron microscope, i.e. it shows a textural autosimilarity regardless the scale. The proposed interpretations therefore should not be extended to the others mineralization hosted in the Neoproterozoic to Cambrian's cover of the Bou Azzer El Graara inlier without any further study. On the other hand, for academic but also applied purposes, this work highlights the necessity to perform more detailed studies of the Cu-occurrences hosted in the cover of the Bou Azzer El Graara inlier to reevaluate the factors controlling these mineralizations.

Acknowledgment

This work was supported by the research project "CALAMINE". Fieldwork has been financed by Managem. Thanks are due to M. Jébrak and P. Ericksson for their accurate and constructive reviews. We are also indebted to I. Di Carlo (ISTO), P. Pehnoud (ISTO), N. Pothier (ISTO), J.-G. Badin (ISTO) and S. Janiec (ISTO) for the sample preparation, the assistance for bibliography, analyses and data treatment.

Bibliography

- 349 Alvaro J.J., Ezzouhairi H., Vennin E., Ribeiro M.L., Clausen S., Charif A., Ait Ayad N., Moreira
350 M.E., 2006. The Early-Cambrian Boho volcano of the El Graara massif, Morocco: Petrology,
351 geodynamic setting and coeval sedimentation. *J. Afr. Earth Sci.* 44, 396-410.
- 352 Benssaou M. et Hamouni N., 1999. Paléoenvironnements et minéralisations de l'Anti-Atlas
353 occidental marocain au Cambrien précocé. *Chron. Rech. Min.* 536-537, 113-119.
- 354 Bouchta R., Boyer F., Routhier P., Saadi M. et Salem M., 1977. L'aire cuprifère de l'Anti-Atlas
355 (Maroc); permanence et arêtes riches. *C. R. Acad. Sc. Paris* 284, 503-506.
- 356 Boudda A., Choubert G., Faure-Muret A., 1979. Essai de stratigraphie de la couverture sédimentaire
357 de l'Anti-Atlas : Adoudounien-Cambrien inférieur. *Notes et Mémoires du Service Géologique du*
358 *Maroc.* 271, 96 pp.
- 359 Choubert G., 1947. L'accident majeur de l'Anti-Atlas. *Comptes Rendus de l'Académie des Sciences*
360 224, 1172-1173.
- 361 Choubert G., 1952. Histoire géologique du domaine de l'Anti-Atlas. In: *Géologie du Maroc. Notes et*
362 *Mémoires du Service Géologique du Maroc* 100, 75-194.
- 363 Compston W., Williams J.L., Kirschvink J.L., Zhang Z., Ma G., 1992. Zircon U-Pb ages for the Early
364 Cambrian time scale. *Journal of the Geological Society, London* 127, 319-332.
- 365 Ducrot J., Lancelot J.R., 1977. Problème de la limite Précambrien-Cambrien: étude
366 radiochronologique par la méthode U/Pb sur zircon du volcan du Jbel Boho. *Can. J. Earth Sci.* 14,
367 1771-1777.
- 368 Dutro, J.T., Dietrich R.V., Foote R.M., 1989. *AGI Data Sheets: or geology in the field,*
369 *laboratory, and office, ed. American Geological Institute, Virginia, U.S.A*

- 370 Faik K., Belfoul M.A., Bouabdelli M. et Hassenforder B., 2002. Les structures de la couverture
371 Néoprotérozoïque terminal et Paléozoïque de la région de Tata, Anti-Atlas centre-occidental,
372 Maroc: déformation polyphasée, ou interactions socle/couverture pendant l'orogénèse
373 hercynienne? *J. African Earth Sci.* 32, 765-776.
- 374 Fossen H., Teyssier C., Whitney D. L., 2013. Transtensional folding. *J. Struct. Geol.* 56, 89-102.
- 375 Jébrak M., 1997. Hydrothermal breccias in vein-type ore deposits: A review of mechanisms,
376 morphology and size distribution. *Ore Geology Reviews* 12, 111-134.
- 377 Kersit, 1984. Internal report, Managem, 40 pp.
- 378 Landing E., 1998. Cambrian subdivisions and correlations. *Can. J. Earth Sci.* 35, 4.
- 379 Leblanc M., 1972. Sur le style disharmonique des plis hercyniens de la couverture, Anti-Atlas central
380 (Maroc). *Comptes Rendus de l'Académie des Sciences* 275, 803-806.
- 381 Leblanc M., 1975. Ophiolites précambriennes et gîtes arsénisés de Cobalt (Bou-Azzer, Maroc). *Notes*
382 *et Mémoires du Service Géologique du Maroc*, 280, 306 pp.
- 383 Leblanc M., 1986. Appareil ignimbrétique et minéralisation cuprifère: Alous (Anti-Atlas, Maroc).
384 *Mineral. Deposita* 21, 129-136.
- 385 Maacha L., Ennaciri A., El Ghorfi M., Baoutoul H., Saquaque A., Soulaïmani A., 2011. The J.
386 La'sal Oxidized Copper Deposit (El Graara inlier, Central Anti-Atlas), in: Mouttaqi A., Rjimati.
387 E.C., Maacha L., Michard A., Soulaïmani A., Ibouh H. (Eds), *New geological and mining*
388 *guidebook of Morocco. Volume 9. Main Mines of Morocco. Service Géologique du Maroc,*
389 *Rabat*, pp. 117-121.
- 390 Maloof A.C., Schrag D.P., Crowley J.L., Bowring S.A., 2005. An expanded record of Early Cambrian
391 carbon cycling from the Anti-Atlas Margin, Morocco. *Can. J. Earth Sci.* 42, 2195-2216.

- 392 Pouit G., 1966. Paléogéographie et répartition des minéralisations stratiformes de cuivre dans l'anti
393 atlas occidental (Maroc). Chron. Rech. Min 356, 279-289.
- 394 Ramdhor P., 1969. The ore minerals and their intergrowths. Pergamon Press, 117 pp.
- 395 Saquaque A., Admou H., Karson J. A., Hefferan, K. & Reuber I., 1989. Precambrian accretionary
396 tectonics in the Bou Azzer–El Graara region, Anti-Atlas, Morocco. *Geology* 17, 1107–1110.
- 397 Sylvester A.G., 1988. Strike-slip faults. *Geol. Soc. Am. Bull.* 100, 1666-1703.
- 398 Skacel J., 1993. Gisement cuprifère polygénétique de Tazalagh (Anti-Atlas occidental). *Mine,*
399 *Géologie & Energie* 54, 127-133.
- 400 Soulaïmani A., Le Corre Cl., Farazdaq R., 1997. Déformation hercynienne et relation
401 socle/couverture dans le domaine du Bas Drâa (Anti-Atlas occidental, Maroc). *J. Afr. Earth Sci.*,
402 24, 271-284.
- 403 Soulaïmani A. and Burkhard M. 2008. The Anti-Atlas chain (Morocco): the southern margin of the
404 Variscan belt along the edge of the West African craton. *Geological Society of London, Special*
405 *Publications* 297, 433-452.
- 406 Soulaïmani A., Egal. E., Razin Ph., Youbi N., Admou H., Blein O., Barbanson L., Gasquet D.,
407 Bouabdelli M., 2013. Notice explicative de la carte géologique du Maroc au 1/50 000, feuille Al
408 Glo'a. Notes et Mémoires du Serv. Géol. Maroc 532 bis, 140 pp.
- 409 Soulaïmanie A., Michard A., Ouanaimi H., Baidder L., Raddi Y., Saddiqi O., Rjimai E.C., 2014. Late
410 Ediacaran-Cambrian structures and their reactivation during the Variscan and Alpine cycles in the
411 Anti-Atlas (Morocco). *J. Afr. Earth Sci.*, 98, 94-112.

Thomas R.J., Fekkak A., Ennih N., Errami E., Loughlin S.C., Gresse P.G., Chevallier L.P., Liégeois J.-P., 2004. A new lithostratigraphic framework for the Anti-Atlas Orogen, Morocco. *J. Afr. Earth Sci.* 39, 217-226.

Figures:

Figure 1: A: Schematic geologic map of the Anti-Atlas with occurrences of copper mineralizations modified from Bouchta et al. (1977). Inliers abbreviations : If, Ifni; Kr, Kerdous; Ir, Igherm; TA, Tagragra d'Akka; TT, Tagragra Tata; AM, Agadir Melloul; Ze, Zenaga; Sr, Sirwa; Bz, Bou Azzer-El Graara; Sg, Saghro; Og, Ougnat. **B:** Geological and structural map of the Bou Azzer-El Graara area with occurrences of copper mineralizations, modified from Leblanc (1975).

Figure 2 : Lithostratigraphic column for the Jbel Laassel area modified from Soulimani et al. (2013).

Figure 3: (a) Geological and structural map of the Jbel Laassel area, modified from Kersit (1984); **(b)** Stereonet (Wulff stereonet, lower hemisphere) of bedding (S_0) and fold axes (n , number of measures).

Figure 4: Geological cross-section (section location A-A' in the map) with ore body limits and mineralogical abundance of the copper mineralization which are represented in a spider diagram. Results are expressed in the corresponding axis of the spider diagram in a scale of 0 to 100% with steps of 20%. Abbreviations: Mal, malachite; Ccl, chrysocolla; Cup, cuprite; Dg, digenite; Bn, bornite; Ccp, chalcopyrite; Cc, chalcocite.

Figure 5: Geological cross-section (section location B-B' in the map) with ore body limits and mineralogical abundance of the copper mineralization which are represented in a spider diagram. Results are expressed in the corresponding axis of the spider diagram in a scale of 0 to 100% with steps of 20%. Same abbreviations than fig.5.

Figure 6: Microphotographs of the Jbel Laassel Cu ore. **A:** Vein showing banding gangue texture with quartz on wall, next dolomite and calcite at center (TL nic. +). **B:** Chalcocite in ribbon between

bands of dolomite and calcite (RL, nic. //). **C:** Chalcocite in vein with hexagonal cleavage (RL, nic. //). **D:** Quartz and chalcocite veinlet link to a vein forming a stockwork (TL, nic. //). **E.1:** Quartz vein cutting bedding (S0) of siltstone and silicification of the siltstones bedding (TL, nic.//). **E.2:** Same photography as E.1 but in RL (nic. //), chalcocite in vein and in dissemination in silicified bedding, these disseminations were observed at SEM scale (see fig.9). **F:** Quartz, dolomite and chalcocite vein showing micro-cockade breccia texture (TL, nic. //). **G:** Quartz in vein with comb texture and malachite veinlets cutting other rock components and structures (TL, nic. +). Abbreviations: Cc1, primary chalcocite; Dol, dolomite; Mal, Malachite; Qtz, quartz. TL, transmitted light; RL, reflected light; nic. //, parallel nicols; nic. +, crossed nicols; S0, bedding of rock.

Figure 7: Paragenetic succession.

Figure 8: Photographs of core drill samples. **A:** Stockwork with gangue minerals, chalcocite, chrysocolla and malachite, and a fluid assisted breccia texture. **B:** Chalcocite stockwork. **C:** Chalcocite stockwork with vein secant and parallel to the dolostone bedding. **D:** Veins of chalcocite cutting the dolostone bedding and chalcocite in dissemination in dolostone but linking to the vein, these disseminations were observed at SEM scale (see fig.9). Abbreviations are given in figure 7.

figure 9: SEM photographs of disseminated copper mineralization (see figures 6E and 7D). **A:** BSE image of a micro-vein (100 μm width) composed by quartz, dolomite and chalcocite. **B:** BSE image of a micro-vein (25 μm width) composed by quartz, dolomite and chalcocite.

Figure 10: Results of PCA analyses of the data base without copper (A, B and C) and of the data base with copper (D and E). **A:** Eigenvalues and cumulative variance function of factors. **B:** Variables projection in the F1-F2 plane (data set without copper: 228 analysis). **C:** samples lithology project in F1-F2 plane (data set without copper: 228 analysis) and Variables projection in the F1-F2 plane without legends. **D:** Variables projection in the F1-F2 plane (data set including copper: 72 analysis). **E:** Variables projection in the F3-F6 plane (data set including copper: 72 analysis).

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8







Figure 9





461

462

463 Highlights:

464

- 465
- 466
- 467
- 468
- Coper mineralization occurs in a Variscan folding band in Cambrian cover.
 - The mineralization appears as a stockwork with autosimilar texture.
 - Results from component principal analyses don't show relation with lithology.
 - The Cu-mineralization is epigenetic, controlled by fold axis.

469